

CHASING SILVER

ANDRZEJ ROSŁANOWSKI AND JURIS STEPRĀNS

ABSTRACT. Answering a question of the first author stated in [6, 0.2] we show that limits of CS iterations of n -Silver forcing notion have the n -localization property.

0. INTRODUCTION

The present paper is concerned with the n -localization property of the n -Silver forcing notion and countable support (CS) iterations of such forcings. The property of n -localization was introduced in Newelski and Rosłanowski [5, p. 826].

Definition 0.1. Let n be an integer greater than 1.

- (1) A tree T is an n -ary tree provided that $(\forall s \in T)(|\text{succ}_T(s)| \leq n)$.
- (2) A forcing notion \mathbb{P} has the n -localization property if

$$\Vdash_{\mathbb{P}} \text{“} (\forall f \in {}^\omega\omega) (\exists T \in \mathbf{V}) (T \text{ is an } n\text{-ary tree and } f \in [T]) \text{”}.$$

Later the n -localization property, the σ -ideal generated by n -ary trees and n -Sacks forcing notion \mathbb{D}_n (see 1.1) have been found applied to problems on convexity numbers of closed subsets of \mathbb{R}^n , see Geschke, Kojman, Kubiś and Schipperus [3], Geschke and Kojman [2] and most recently Geschke [1].

We do not have any result of the form “CS iteration of proper forcing notions with the n -localization property has the n -localization” yet. A somewhat uniform and general treatment of preserving the n -localization has been recently presented in [6]. However, the treatment in that paper does not cover the n -Silver forcing notion \mathbb{S}_n (see 1.1), as a matter of fact it was not clear at some moment if \mathbb{S}_n has the property at all. It was stated in [5, Theorem 2.3] that the same proof as for \mathbb{D}_n works also for CS iterations and products of the n -Silver forcing notions \mathbb{S}_n (see Definition 1.1(3)). Maybe some old wisdom got lost, but it does not look like that *the same arguments work for the n -Silver forcing \mathbb{S}_n* . In the present paper we correct this gap and we provide a full proof that CS iterations of \mathbb{S}_n (and other forcings listed in 1.1) have the n -localization property, see 1.6. Our main result 1.5 on the n -Silver forcing seems to be very \mathbb{S}_n -specific and it is not clear to which extend it may be generalized.

Let us explain what is a possible problem with the n -Silver forcing – let us look at the “classical” Silver forcing \mathbb{S}_2 . Given a Silver condition f such that $f \Vdash_{\mathbb{S}_2} \dot{\tau} \in {}^\omega\omega$, standard arguments allow it to be assumed that the complement of the domain of f can be enumerated in the increasing order as $\{k_i : i < \omega\}$ and that for each

Date: August 2005.

1991 Mathematics Subject Classification. 03E40, 03E35.

Key words and phrases. n -localization property, the Silver forcing, CS iterations.

The research of the second author was partially supported by NSERC of Canada.

$i \in \omega$ and $\rho : \{k_j : j < i\} \rightarrow 2$ the condition $f \cup \rho$ decides the value of $\mathcal{T} \upharpoonright i$, say $f \cup \rho \Vdash \mathcal{T} \upharpoonright i = \sigma_\rho$. Now one could take the tree

$$T^\oplus = \{\nu \in {}^\omega \omega : (\exists i < \omega)(\exists \rho \in \{k_j : j < i\} 2)(\nu \leq \sigma_\rho)\}.$$

Easily $p \Vdash \mathcal{T} \in [T^\oplus]$, but T^\oplus does not have to be a binary tree! (It could well be that $\sigma_\rho = \sigma^*$ for all ρ of length 100 and then $\sigma_{\rho'}$ for ρ' of length 101 are pairwise distinct.) So we would like to make sure that σ_ρ for ρ 's of the same length are distinct, but this does not have to be possible. To show that \mathbb{S}_2 has the 2-localization property we have to be a little bit more careful. Let us give a combinatorial result which easily implies that \mathbb{S}_2 has the 2-localization property. Its proof is in the heart of our proof of Theorem 1.5.

Fix $\Psi : {}^\omega 2 \rightarrow \omega$. Define $\Psi^* : {}^\omega 2 \rightarrow {}^\omega \omega$ by induction: Let $\Psi^*(\langle \rangle) = \langle \rangle$ and define $\Psi^*(t \frown \langle i \rangle) = \Psi^*(t) \frown \langle \Psi(t \frown \langle i \rangle) \rangle$. If ξ is a partial function from ω to 2 and $\ell \leq \omega$ define

$$W^\ell(\xi) = \{t \in {}^{m 2} : m < \min(\ell + 1, \omega) \ \& \ \xi \upharpoonright m \subseteq t\}$$

and then define $T^\ell(\xi) = \{\Psi^*(t) : t \in W^\ell(\xi)\}$, $T(\xi) = T^\omega(\xi)$.

Theorem 0.2. *For any $\Psi : {}^\omega 2 \rightarrow \omega$ there is a partial function $\xi : \omega \rightarrow 2$ with co-infinite domain such that $T(\xi)$ is a binary tree.*

Proof. To begin, two equivalence relations on ${}^\omega 2$ will be defined. First, define $s \equiv t$ if and only if $\Psi(t \frown \theta) = \Psi(s \frown \theta)$ for all $\theta \in {}^\omega 2$. Next, define $s \sim t$ if and only if $\Psi^*(s) = \Psi^*(t)$.

Now construct by induction on $m < \omega$ an increasing sequence

$$x_0 < x_1 < \dots < x_m < N_m$$

and $\xi_m : N_m \setminus \{x_0, x_1, \dots, x_m\} \rightarrow 2$ such that $T^{N_m}(\xi_m)$ is a binary branching tree and, moreover, if s and t are maximal elements of $W^{N_m}(\xi_m)$ and $t \sim s$ then $t \equiv s$. The induction starts with $N_0 = 0$. If the induction has been completed for m then let $x_{m+1} = N_m$. Let $\Delta = \{d_0, d_1, \dots, d_j\}$ be a set of maximal elements of $T^{N_m}(\xi_m)$ such that precisely one member of each \sim equivalence class belongs to Δ . Now, by induction on $i \leq j$ define N^i and $\xi^i : N^i \setminus (N_m + 1) \rightarrow 2$ as follows. Let $N^0 = N_m + 1$ and let $\xi^0 = \emptyset$. Given N^i and ξ^i , if there is some $N > N^i$ and $\xi \supseteq \xi^i$ such that $d_i \frown \langle 0 \rangle \frown \xi \equiv d_i \frown \langle 1 \rangle \frown \xi$ then let $N^{i+1} = N$ and let $\xi^{i+1} = \xi$. Otherwise it must be the case that $d_i \frown \langle 0 \rangle \frown \xi^i \not\equiv d_i \frown \langle 1 \rangle \frown \xi^i$ and so it must be possible to find $N^{i+1} > N^i$ and $\xi^{i+1} \supseteq \xi^i$ such that $d_i \frown \langle 0 \rangle \frown \xi^{i+1} \not\sim d_i \frown \langle 1 \rangle \frown \xi^{i+1}$. Finally, let $N_{m+1} = N^j$ and $\xi_{m+1} = \xi_m \cup \xi^j$.

To see that this works, it must be shown that $T^{N_{m+1}}(\xi_{m+1})$ is a binary tree and that if s and t are maximal elements of $W^{N_{m+1}}(\xi_{m+1})$ and $t \sim s$ then $t \equiv s$. To check the first condition it suffices to take t a maximal element of $T^{N_m}(\xi_m)$ and check that the tree $T^{N_{m+1}}(\xi_{m+1})$ above t is binary. Then $t = \Psi^*(d_i)$ for some i by the induction hypothesis. The tree $T^{N_{m+1}}(\xi_{m+1})$ above t consists is generated by all $\Psi^*(d \frown \langle a \rangle \frown \xi_j)$ where $d \sim d_i$ and $a \in 2$. Note however that if $d \sim d_i$ then $d \equiv d_i$ and so

$$\Psi^*(d \frown \langle a \rangle \frown \xi_j) = \Psi^*(d_i \frown \langle a \rangle \frown \xi_j).$$

Therefore $\Psi^*(d \frown \langle a \rangle \frown \xi_j)$ depends only on a and not on d and so $T^{N_{m+1}}(\xi_{m+1})$ is binary above t .

To check the second condition suppose that s and t are maximal elements of $W^{N_{m+1}}(\xi_{m+1})$ and $t \sim s$. This implies that $t \upharpoonright N_m \sim s \upharpoonright N_m$ and hence $t \upharpoonright N_m \equiv$

$s \upharpoonright N_m$. Let i be such that $t \upharpoonright N_m \sim s \upharpoonright N_m \sim d_i$. If $t(N_m) = s(N_m) = y$ then $t = t \upharpoonright N_m \frown \langle y \rangle \frown \xi^j$ and $s = s \upharpoonright N_m \frown \langle y \rangle \frown \xi^j$ and, since $t \upharpoonright N_m \equiv s \upharpoonright N_m$, it is immediate that $t \equiv s$. So assume that $t(N_m) = 0$ and $s(N_m) = 1$. By the same argument it follows that $t \equiv d_i \frown \langle 0 \rangle \frown \xi^j$ and $s \equiv d_i \frown \langle 1 \rangle \frown \xi^j$. Hence it suffices to show that $d_i \frown \langle 0 \rangle \frown \xi^j \equiv d_i \frown \langle 1 \rangle \frown \xi^j$. Note that $d_i \frown \langle 0 \rangle \frown \xi^j \sim d_i \frown \langle 1 \rangle \frown \xi^j$ since $t \sim d_i \frown \langle 0 \rangle \frown \xi^j$ and $s \sim d_i \frown \langle 1 \rangle \frown \xi^j$. This means that it must have been possible to find ξ^i such that $d_i \frown \langle 0 \rangle \frown \xi^i \equiv d_i \frown \langle 1 \rangle \frown \xi^i$. It follows that $d_i \frown \langle 0 \rangle \frown \xi^j \equiv d_i \frown \langle 1 \rangle \frown \xi^j$. \square

Notation: Our notation is rather standard and compatible with that of classical textbooks (like Jech [4]). In forcing we keep the older convention that *a stronger condition is the larger one*.

- (1) n is our fixed integer, $n \geq 2$.
- (2) For two sequences η, ν we write $\nu \triangleleft \eta$ whenever ν is a proper initial segment of η , and $\nu \trianglelefteq \eta$ when either $\nu \triangleleft \eta$ or $\nu = \eta$. The length of a sequence η is denoted by $\text{lh}(\eta)$.
- (3) A *tree* is a family of finite sequences closed under initial segments. For a tree T and $\eta \in T$ we define *the successors of η in T* and *maximal points of T* by:

$$\begin{aligned} \text{succ}_T(\eta) &= \{ \nu \in T : \eta \triangleleft \nu \ \& \ \neg(\exists \rho \in T)(\eta \triangleleft \rho \triangleleft \nu) \}, \\ \text{max}(T) &= \{ \nu \in T : \text{there is no } \rho \in T \text{ such that } \nu \triangleleft \rho \}. \end{aligned}$$

For a tree T the family of all ω -branches through T is denoted by $[T]$.

- (4) For a forcing notion \mathbb{P} , all \mathbb{P} -names for objects in the extension via \mathbb{P} will be denoted with a tilde below (e.g., τ, X).

1. DEFINITIONS AND THE RESULT

Definition 1.1. (1) *The n -Sacks forcing notion \mathbb{D}_n* consists of perfect trees $p \subseteq {}^\omega n$ such that

$$(\forall \eta \in p)(\exists \nu \in p)(\eta \triangleleft \nu \ \& \ \text{succ}_p(\eta) = n).$$

The order of \mathbb{D}_n is the reverse inclusion, i.e., $p \leq_{\mathbb{D}_n} q$ if and only if $q \subseteq p$. (See [5].)

- (2) *The uniform n -Sacks forcing notion \mathbb{Q}_n* consists of perfect trees $p \subseteq {}^\omega n$ such that

$$(\exists X \in [\omega]^\omega)(\forall \eta \in p)(\text{lh}(\eta) \in X \Rightarrow \text{succ}_p(\eta) = n).$$

The order of \mathbb{Q}_n is the reverse inclusion, i.e., $p \leq_{\mathbb{Q}_n} q$ if and only if $q \subseteq p$. (See [7].)

- (3) Let us assume that $G = (V, E)$ is a hypergraph on a Polish space V such that

- $E \subseteq [V]^{n+1}$ is open in the topology inherited from V^{n+1} , and
- $(\forall e \in E)(\forall v \in V \setminus e)(\exists w \in e)((e \setminus \{w\}) \cup \{v\} \in E)$,
- for every non-empty open subset U of V and every countable family \mathcal{F} of subsets of U , either $\bigcup \mathcal{F} \neq U$ or $[F]^{n+1} \cap E \neq \emptyset$ for some $F \in \mathcal{F}$.

The Geschke forcing notion \mathbb{P}_G for G consists of all closed sets $C \subseteq V$ such that the hypergraph $(C, E \cap [C]^{n+1})$ is uncountably chromatic on every non-empty open subset of C . The order of \mathbb{P}_G is the inverse inclusion, i.e., $C \leq_{\mathbb{P}_G} D$ if and only if $D \subseteq C$. (See [1].)

- Definition 1.2.** (1) The n -Silver forcing notion \mathbb{S}_n consists of partial functions f such that $\text{Dom}(f) \subseteq \omega$, $\text{Rng}(f) \subseteq n$ and $\omega \setminus \text{Dom}(f)$ is infinite. The order of \mathbb{S}_n is the inclusion, i.e., $f \leq_{\mathbb{S}_n} g$ if and only if $f \subseteq g$.
- (2) For an integer $i \in \omega$ and a condition $f \in \mathbb{S}_n$ we let $\text{FP}_i(f)$ to be the unique element of $\omega \setminus \text{Dom}(f)$ such that $|\text{FP}_i(f) \setminus \text{Dom}(f)| = i$. (The FP stands for *Free Point*.)
- (3) A binary relation \leq_i^* on \mathbb{S}_n is defined by $f \leq_i^* g$ if and only if $(f, g \in \mathbb{S}_n)$ and $f \leq_{\mathbb{S}_n} g$ and

$$(\forall j \in \omega)(j < \lfloor i/4 \rfloor \Rightarrow \text{FP}_j(f) = \text{FP}_j(g)).$$

- (4) For $f \in \mathbb{S}_n$ and $\sigma : N \rightarrow n$, $N < \omega$ we define $f * \sigma$ as the unique condition in \mathbb{S}_n such that $\text{Dom}(f * \sigma) = \text{Dom}(f) \cup \{\text{FP}_i(f) : i < N\}$, $f \subseteq f * \sigma$ and $f * \sigma(\text{FP}_i(f)) = \sigma(i)$ for $i < N$.

Definition 1.3. Let \mathbb{P} be a forcing notion.

- (1) For a condition $p \in \mathbb{P}$ we define a game $\mathcal{D}_n^\ominus(p, \mathbb{P})$ of two players, *Generic* (“she”) and *Antigeneric* (“he”). A play of $\mathcal{D}_n^\ominus(p, \mathbb{P})$ lasts ω moves and during it the players construct a sequence $\langle (s_i, \bar{\eta}^i, \bar{p}^i, \bar{q}^i) : i < \omega \rangle$ as follows. At a stage $i < \omega$ of the play, first Generic chooses a finite n -ary tree s_i such that

$$(\alpha) \quad |\max(s_0)| \leq n \text{ and if } i = j + 1 \text{ then } s_j \text{ is a subtree of } s_i \text{ such that}$$

$$(\forall \eta \in \max(s_i))(\exists \ell < \text{lh}(\eta))(\eta \upharpoonright \ell \in \max(s_j)),$$

and

$$(\forall \nu \in \max(s_j))(0 < |\{\eta \in \max(s_i) : \nu \triangleleft \eta\}| \leq n).$$

Next

(β) Generic picks an enumeration $\bar{\eta}^i = \langle \eta_\ell^i : \ell < k_i \rangle$ of $\max(s_i)$ (so $k_i < \omega$) and then the two players play a subgame of length k_i choosing successive terms of a sequence $\langle p_{\eta_\ell^i}^i, q_{\eta_\ell^i}^i : \ell < k_i \rangle$. At a stage $\ell < k_i$ of the subgame, first Generic picks a condition $p_{\eta_\ell^i}^i \in \mathbb{P}$ such that

$$(\gamma)_\ell^i \text{ if } j < i, \nu \in \max(s_j) \text{ and } \nu \triangleleft \eta_\ell^i, \text{ then } q_\nu^j \leq p_{\eta_\ell^i}^i \text{ and } p \leq p_{\eta_\ell^i}^i,$$

and then Antigeneric answers with a condition $q_{\eta_\ell^i}^i$ stronger than $p_{\eta_\ell^i}^i$.

Finally, Generic wins the play $\langle (s_i, \bar{\eta}^i, \bar{p}^i, \bar{q}^i) : i < \omega \rangle$ if and only if

$$(\otimes) \text{ there is a condition } q \geq p \text{ such that for every } i < \omega \text{ the family } \{q_\eta^i : \eta \in \max(s_i)\} \text{ is predense above } q.$$

- (2) We say that \mathbb{P} has the \ominus_n -property whenever Generic has a winning strategy in the game $\mathcal{D}_n^\ominus(p, \mathbb{P})$ for any $p \in \mathbb{P}$.
- (3) Let $K \in [\omega]^\omega$, $p \in \mathbb{P}$. A strategy **st** for Generic in $\mathcal{D}_n^\ominus(p, \mathbb{P})$ is K -nice whenever
- ($\boxtimes_{\text{nice}}^K$) if so far Generic used **st** and s_i and $\bar{\eta}^i = \langle \eta_\ell^i : \ell < k \rangle$ are given to her as innings at a stage $i < \omega$, then
- $s_i \subseteq \bigcup_{j \leq i+1} s_j$, $\max(s_i) \subseteq {}^{(i+1)}(n+1)$ and
 - if $\eta \in \max(s_i)$ and $i \notin K$, then $\eta(i) = n$, and
 - if $\eta \in \max(s_i)$ and $i \in K$, then $\text{succ}_{s_i}(\eta \upharpoonright i) = n$,

- if $i \in K$ and $\langle p_{\eta_i}^i, q_{\eta_i}^i : \ell < k \rangle$ is the result of the subgame of level i in which Generic uses **st**, then the conditions $p_{\eta_i}^i$ (for $\ell < k$) are pairwise incompatible.
- (4) We say that \mathbb{P} has the *nice* \ominus_n -property if for every $K \in [\omega]^\omega$ and $p \in \mathbb{P}$, Generic has a K -nice winning strategy in $\mathcal{D}_n^\ominus(p, \mathbb{P})$.

Theorem 1.4 (See [6, 3.1+1.6+1.4]). *The limits of CS iterations of the forcing notions defined in 1.1, 1.2 have the nice \ominus_n -property.*

Now we may formulate our main result.

Theorem 1.5. *Assume that \mathbb{P} has the nice \ominus_n -property and the n -localization property. Let \mathbb{S}_n be the \mathbb{P} -name for the n -Silver forcing notion. Then the composition $\mathbb{P} * \mathbb{S}_n$ has the n -localization property.*

The proof of Theorem 1.5 is presented in the following section. Let us note here that this theorem implies n -localization for CS iterations of the forcing notions mentioned here.

Corollary 1.6. *Let $\mathbb{Q} = \langle \mathbb{P}_\xi, \mathbb{Q}_\xi : \xi < \gamma \rangle$ be a CS iteration such that, for every $\xi < \gamma$, \mathbb{Q}_ξ is a \mathbb{P}_ξ -name for one of the forcing notions defined in 1.1, 1.2. Then $\mathbb{P}_\gamma = \lim(\mathbb{Q})$ has the n -localization property.*

Proof. By induction on γ .

If $\gamma = \gamma_0 + 1$ and \mathbb{Q}_{γ_0} is a \mathbb{P}_{γ_0} -name for the n -Silver forcing notion, then 1.5 applies. (Note that \mathbb{P}_{γ_0} has the nice \ominus_n -property by 1.4 and it has the n -localization property by the inductive hypothesis.)

If $\gamma = \gamma_0 + 1$ and \mathbb{Q}_{γ_0} is a \mathbb{P}_{γ_0} -name for \mathbb{D}_n or \mathbb{Q}_n or \mathbb{P}_G , then [6, Theorem 3.4] applies. (Note that \mathbb{P}_{γ_0} has the nice \ominus_n -property by 1.4 and it has the n -localization property by the inductive hypothesis.)

If γ is limit then [6, 3.5] applies. □

Corollary 1.7. *No CS iteration of \mathbb{S}_2 adds an \mathbb{S}_4 -generic real.*

2. PROOF OF THEOREM 1.5

Let τ be a $\mathbb{P} * \mathbb{S}_n$ -name for a member of ${}^\omega\omega$. We may assume that $\Vdash_{\mathbb{P} * \mathbb{S}_n} \tau \notin \mathbf{V}$. If $G \subseteq \mathbb{P}$ is generic over \mathbf{V} , then we will use the same notation τ for \mathbb{S}_n -name in $\mathbf{V}[G]$ for a member of ${}^\omega\omega$ that is given by the original τ in the extension via $\mathbb{P} * \mathbb{S}_n$.

Let $(p, \underline{f}) \in \mathbb{P} * \mathbb{S}_n$ and let **st** be a winning strategy of Generic in $\mathcal{D}_n^\ominus(p, \mathbb{P})$ which is nice for the set $K = \{4j + 2 : j \in \omega\}$ (see 1.3(3)).

By induction on i we are going to choose for each $i < \omega$

$$s_i, \bar{\eta}^i, \bar{p}^i, \bar{q}^i, \underline{f}_i,$$

and for also $m_i, \bar{\sigma}^i$ for odd $i < \omega$ such that the following conditions $(\boxtimes)_1 - (\boxtimes)_7$ are satisfied.

- (\boxtimes)₁ $\langle s_i, \bar{\eta}^i, \bar{p}^i, \bar{q}^i : i < \omega \rangle$ is a play of $\mathcal{D}_n^\ominus(p, \mathbb{P})$ in which Generic uses **st**.
- (\boxtimes)₂ \underline{f}_i is a \mathbb{P} -name for a condition in \mathbb{S}_n , and we stipulate that $\underline{f}_{-1} = \underline{f}$.
- (\boxtimes)₃ $q_\eta^i \Vdash_{\mathbb{P}} \underline{f}_{i-1} \leq_i^* \underline{f}_i$ for each $\eta \in \max(s_i)$.

For odd $i < \omega$:

- (\boxtimes)₄ $m_i < m_{i+2} < \omega$, $\bar{\sigma}^i = \langle \sigma_{\rho, \eta}^i : \eta \in \max(s_i) \ \& \ \rho \in {}^{[i/4]}n \rangle$, $\sigma_{\rho, \eta}^i : m_i \rightarrow \omega$.
- (\boxtimes)₅ $(q_\eta^i, \underline{f}_i * \rho) \Vdash_{\mathbb{P} * \mathbb{S}_n} \tau \upharpoonright m_i = \sigma_{\rho, \eta}^i$ for $\rho \in {}^{[i/4]}n$ and $\eta \in \max(s_i)$.

(\boxtimes)₆ If $\eta \in \max(s_i)$ and $\rho, \rho' : [i/4] \rightarrow n$ are distinct but $\sigma_{\rho, \eta}^i = \sigma_{\rho', \eta}^i$, then for every $q \geq q_\eta^i$ and a \mathbb{P} -name \underline{g} for an n -Silver condition and m, σ, σ' such that

$$q \Vdash_{\mathbb{P}} \underline{f}_i \leq_i^* \underline{g}, \quad (q, \underline{g} * \rho) \Vdash_{\mathbb{P} * \mathbb{S}_n} \tau \upharpoonright m = \sigma, \quad (q, \underline{g} * \rho') \Vdash_{\mathbb{P} * \mathbb{S}_n} \tau \upharpoonright m = \sigma'$$

we have $\sigma = \sigma'$.

(\boxtimes)₇ If $\eta, \eta' \in \max(s_i)$ are distinct, $\rho, \rho' : [i/4] \rightarrow n$, then $\sigma_{\rho, \eta}^i \neq \sigma_{\rho', \eta'}^i$.

So suppose that $i < \omega$ is even and we have already defined $s_{i-1}, \bar{q}^{i-1}, m_{i-1}$ and \underline{f}_{i-1} (we stipulate $s_{-1} = \{\langle \rangle\}$, $q_{\langle \rangle}^{-1} = p$, $\underline{f}_{-1} = \underline{f}$ and $m_{-1} = 0$). Let $j = \lfloor i/4 \rfloor$ (so either $i = 4j$ or $i = 4j + 2$).

The strategy **st** and demand (\boxtimes)₁ determine s_i and $\bar{\eta}^i = \langle \eta_k^i : k < k_i \rangle$. To define \bar{p}^i, \bar{q}^i and \underline{f}_i we consider the following run of the subgame of level i of $\mathfrak{D}_n^\ominus(p, \mathbb{P})$. Assume we are at stage $k < k_i$ of the subgame. Now, $p_{\eta_k^i}^i$ is given by the strategy **st** (and (\boxtimes)₁, of course). Suppose for a moment that $G \subseteq \mathbb{P}$ is generic over \mathbf{V} , $p_{\eta_k^i}^i \in G$. Working in $\mathbf{V}[G]$ we may choose $\bar{\ell}, \bar{L}, g^*, \bar{\sigma}^*, M$ such that

- (\boxtimes)₈ ^{α} $M = n^j$, $\bar{\ell} = \langle \ell_m : m \leq M \rangle$ and $j = \ell_0 < \dots < \ell_M$, $\bar{L} = \langle L_m : m \leq M \rangle$ and $m_{i-1} < L_0 < \dots < L_M$,
- (\boxtimes)₈ ^{β} $g^* \in \mathbb{S}_n$, $\underline{f}_{i-1}[G] \leq_i^* g^*$ and $\bar{\sigma}^* = \langle \sigma_\rho^* : \rho \in {}^{\ell_M} n \rangle$, $\sigma_\rho^* \in {}^{L_M} \omega$ (for $\rho \in {}^{\ell_M} n$),
- (\boxtimes)₈ ^{γ} $g^* * (\rho \upharpoonright \ell_m) \Vdash_{\mathbb{S}_n} \tau \upharpoonright L_m = \sigma_\rho^* \upharpoonright L_m$ for each $m \leq M$ and $\rho \in {}^{\ell_M} n$,
- (\boxtimes)₈ ^{δ} if $\rho_0, \rho_1 \in {}^{\ell_M} n$, $\rho_0 \upharpoonright j \neq \rho_1 \upharpoonright j$ but $\sigma_{\rho_0}^* \upharpoonright L_0 = \sigma_{\rho_1}^* \upharpoonright L_0$, then there is no condition $g \in \mathbb{S}_n$ such that $g^* \leq_i^* g$ and for some $L < \omega$ and distinct $\sigma_0, \sigma_1 \in {}^L \omega$ we have that $g * \rho_0 \Vdash \tau \upharpoonright L = \sigma_0$, $g * \rho_1 \Vdash \tau \upharpoonright L = \sigma_1$,
- (\boxtimes)₈ ^{ε} for each $m < M$ and $\rho_0 \in {}^{\ell_m} n$ the set $\{\sigma_\rho^* \upharpoonright [L_m, L_{m+1}) : \rho_0 \triangleleft \rho \in {}^{\ell_m} n\}$ has at least $n^j \cdot k_i + 777$ elements.

It should be clear how the construction is done. (First we take care of clause (\boxtimes)₈ ^{δ} by going successively through all pairs of elements of ${}^j n$ and trying to force distinct values for initial segments of τ , if only this is possible. Then we ensure (\boxtimes)₈ ^{ε} basically by deciding longer and longer initial segments of τ on fronts/levels of a fusion sequence of conditions in \mathbb{S}_n and using the assumption that τ is forced to be “new”.) Now, going back to \mathbf{V} , we may choose a condition $q_{\eta_k^i}^i \in \mathbb{P}$ stronger than $p_{\eta_k^i}^i$ and a \mathbb{P} -name $\underline{g}^{*,k}$ for a condition in \mathbb{S}_n and objects $\bar{\ell}^k, \bar{L}^k, \bar{\sigma}^{*,k}$ such that

$$q_{\eta_k^i}^i \Vdash_{\mathbb{P}} \text{“ } \bar{\ell}^k, \bar{L}^k, \underline{g}^{*,k}, \bar{\sigma}^{*,k}, n^j \text{ satisfy clauses } (\boxtimes)_{8}^{\alpha} - (\boxtimes)_{8}^{\varepsilon} \text{ as } \bar{\ell}, \bar{L}, g^*, \bar{\sigma}^*, M \text{ there ”.}$$

The condition $q_{\eta_k^i}^i$ is treated as an inning of Antigeneric at stage k of the subgame of $\mathfrak{D}_n^\ominus(p, \mathbb{P})$ and the process continues.

After the subgame of level i is completed, we have defined \bar{p}^i and \bar{q}^i . We also choose \underline{f}_i to be a \mathbb{P} -name for an element of \mathbb{S}_n such that $\Vdash_{\mathbb{P}} \underline{f}_{i-1} \leq_i^* \underline{f}_i$ and $q_{\eta_k^i}^i \Vdash_{\mathbb{P}} \underline{f}_i = \underline{g}^{*,k}$ for all $k < k_i$ (remember that **st** is nice, so the conditions $q_{\eta_k^i}^i$ are pairwise incompatible). This completes the description of what happens at the stage i of the construction (one easily verifies that (\boxtimes)₁–(\boxtimes)₃ are satisfied) and we proceed to the next, $i + 1$, stage. Note that $\lfloor (i + 1)/4 \rfloor = j$.

We let $m_{i+1} = \max(L_M^k : k < k_i) + 5$ and let $\ell = \max(\ell_M^k : k < k_i) + 5$. Similarly as at stage i , s_{i+1} and $\bar{\eta}^{i+1} = \langle \eta_k^{i+1} : k < k_{i+1} \rangle$ are determined by the strategy **st** and (\boxtimes)₁; note that $\max(s_{i+1}) = \{\nu \frown \langle n \rangle : \nu \in \max(s_i)\}$ so $k_{i+1} = k_i$. To define

$\bar{p}^{i+1}, \bar{q}^{i+1}$ and \underline{f}_{i+1} we consider the following round of the subgame of level $i+1$ of $\mathcal{D}_n^\ominus(p, \mathbb{P})$. At a stage $k < k_{i+1}$ of the subgame, letting $\eta = \eta_k^{i+1}$, the condition p_η^{i+1} is given by the strategy \mathbf{st} . Suppose for a moment that $G \subseteq \mathbb{P}$ is generic over \mathbf{V} , $p_\eta^{i+1} \in G$. In $\mathbf{V}[G]$ we may choose a condition $h^* \in \mathbb{S}_n$ such that

(\boxtimes)₉ $\underline{f}_i[G] \leq_\ell^* h^*$ and for every $\rho \in {}^\ell n$ the condition $h^* * \rho$ decides the value of $\mathcal{I} \upharpoonright m_{i+1}$, say $h^* * \rho \Vdash_{\mathbb{S}_n} \mathcal{I} \upharpoonright m_{i+1} = \sigma_\rho$.

Then going back to \mathbf{V} we choose a \mathbb{P} -name $\underline{h}^{*,\eta}$ for a condition in \mathbb{S}_n , a sequence $\bar{\sigma}^\eta = \langle \sigma_\rho^\eta : \rho \in {}^\ell n \rangle$ and a condition $q_\eta^{i+1} \geq p_\eta^{i+1}$ such that

$$q_\eta^{i+1} \Vdash_{\mathbb{P}} \text{“ } \underline{h}^{*,\eta}, \bar{\sigma}^\eta \text{ are as in } (\boxtimes)_9 \text{”}.$$

The condition q_η^{i+1} is treated as an inning of Antigeneric at stage k of the subgame of $\mathcal{D}_n^\ominus(p, \mathbb{P})$ and the process continues.

After the subgame of level $i+1$ is completed, we have defined \bar{p}^{i+1} and \bar{q}^{i+1} . Since for every $\eta \in \max(s_{i+1})$ we have that $p_\eta^{i+1} \geq q_\eta^{i+1}$, we may use (\boxtimes)₈ and choose $\rho(\eta) : [j, \ell] \rightarrow n$ (for $\eta \in \max(s_{i+1})$) such that

(\boxtimes)₁₀ if $\eta, \eta' \in \max(s_{i+1})$ are distinct and $\theta, \theta' \in {}^j n$, and $\rho = \theta \frown \rho(\eta)$, $\rho' = \theta' \frown \rho(\eta')$, then $\sigma_\rho^\eta \neq \sigma_{\rho'}^{\eta'}$.

Let \underline{f}_i be a \mathbb{P} -name for a condition in \mathbb{S}_n such that $\Vdash_{\mathbb{P}} \underline{f}_i \leq_{i+1}^* \underline{f}_{i+1}$ and

$$q_\eta^{i+1} \Vdash_{\mathbb{P}} \text{“ } \underline{h}^{*,\eta} \leq_i^* \underline{f}_i \ \& \ (\forall \theta \in {}^j n) (\underline{f}_i * \theta = \underline{h}^{*,\eta} * (\theta \frown \rho(\eta))) \text{”}.$$

Also, for $\eta \in \max(s_{i+1})$ and $\rho \in {}^j n$, we let $\sigma_{\rho,\eta}^{i+1} = \sigma_{\rho \frown \rho(\eta)}^\eta$. This completes the description of what happens at the stage $i+1$ of the construction (one easily checks that (\boxtimes)₁–(\boxtimes)₇ are satisfied). Thus we have finished the description of the inductive step of the construction of $s_i, \bar{\eta}^i, \bar{p}^i, \bar{q}^i, \underline{f}_i$ (for $i < \omega$).

After the construction is carried out we may pick a condition $q \in \mathbb{P}$ stronger than p and such that for each $i < \omega$ the family $\{q_\eta^i : \eta \in \max(s_i)\}$ is predense above q (possible by (\boxtimes)₁).

Suppose that $G \subseteq \mathbb{P}$ is generic over \mathbf{V} , $q \in G$. Then there is $\eta \in {}^\omega(n+1)$ such that $\eta \upharpoonright (i+1) \in \max(s_i)$ and $q_{\eta \upharpoonright (i+1)}^i \in G$ for each $i < \omega$. Therefore we may use (\boxtimes)₃ to conclude that there is a condition $g \in \mathbb{S}_n$ stronger than all $\underline{f}_i[G]$. Going back to \mathbf{V} , we may choose a \mathbb{P} -name \underline{g} for a condition in \mathbb{S}_n such that $q \Vdash_{\mathbb{P}} (\forall i < \omega) (\underline{f}_i \leq \underline{g})$.

Note that for each $i < \omega$ the family $\{(q_\eta^i, \underline{f}_i * \rho) : \eta \in \max(s_i) \ \& \ \rho \in {}^{[i/4]}n\}$ is predense in $\mathbb{P} * \mathbb{S}_n$ above (q, \underline{g}) , and hence (by (\boxtimes)₅)

$$(q, \underline{g}) \Vdash_{\mathbb{P} * \mathbb{S}_n} \text{“ } \mathcal{I} \upharpoonright m_i \in \{\sigma_{\rho,\eta}^i : \eta \in \max(s_i) \ \& \ \rho \in {}^{[i/4]}n\} \text{ for every odd } i < \omega \text{”}.$$

Also,

(\boxtimes)₁₁ if $i \geq 3$ is odd, $\eta \in \max(s_i)$, $\rho \in {}^{[i/4]}n$ and $\eta' = \eta \upharpoonright (i-1)$ and $\rho' = \rho \upharpoonright [(i-2)/4]$, then $\eta' \in \max(s_{i-2})$ and $\sigma_{\rho',\eta'}^{i-2} = \sigma_{\rho,\eta}^i \upharpoonright m_{i-2}$.

[Why? Since \mathbf{st} is a nice strategy, $\eta \upharpoonright i \in \max(s_{i-1})$ and $\eta' \in \max(s_{i-2})$. It follows from (\boxtimes)₁ that $q_{\eta'}^{i-2} \leq q_{\eta \upharpoonright i}^{i-1} \leq q_\eta^i$ and by (\boxtimes)₃ we have $q_\eta^i \Vdash_{\mathbb{P}} \underline{f}_{i-2} \leq_{i-1}^* \underline{f}_i$. Therefore $q_\eta^i \Vdash_{\mathbb{P}} \underline{f}_{i-2} * \rho' \leq \underline{f}_i * \rho$ and $(q_{\eta'}^{i-2}, \underline{f}_{i-2} * \rho') \leq (q_\eta^i, \underline{f}_i * \rho)$, so using (\boxtimes)₅ we may conclude that $\sigma_{\rho',\eta'}^{i-2} = \sigma_{\rho,\eta}^i \upharpoonright m_{i-2}$.]

Let

$$T = \{\nu \in {}^\omega > \omega : (\exists i < \omega \text{ odd}) (\exists \eta \in \max(s_i)) (\exists \rho \in {}^{[i/4]}n) (\nu \leq \sigma_{\rho,\eta}^i)\}.$$

Then T is a perfect tree and $(q, g) \Vdash_{\mathbb{P} * \mathbb{S}_n} \tau \in [T]$. So the theorem will readily follow once we show that T is n -ary. To this end we are going to argue that

(\boxtimes)₁₂ if $i \geq 3$ is odd, $\eta \in \max(s_i)$, $\rho \in {}^{[i/4]}n$, then

$$\left| \left\{ \sigma_{\pi, \nu}^i : \nu \in \max(s_i) \ \& \ \pi \in {}^{[i/4]}n \ \& \ \sigma_{\rho, \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \nu}^i \upharpoonright m_{i-2} \right\} \right| \leq n.$$

CASE A: $i = 4j + 1$ for some $j < \omega$.

Suppose that $\eta, \nu \in \max(s_i)$, $\rho, \pi \in {}^{[i/4]}n$ are such that $\sigma_{\rho, \eta}^i \neq \sigma_{\pi, \nu}^i$ but $\sigma_{\rho, \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \nu}^i \upharpoonright m_{i-2}$. The latter and (\boxtimes)₇ imply that $\eta \upharpoonright (i-1) = \nu \upharpoonright (i-1)$, and since $i-1, i \notin K$ we get that $\eta(i-1) = \nu(i-1) = n = \eta(i) = \nu(i)$ (remember: **st** is nice for K), so $\eta = \nu$. If $\rho \upharpoonright (j-1) \neq \pi \upharpoonright (j-1)$, then let $\rho' = \rho \upharpoonright (j-1) \frown (\pi \upharpoonright (j-1))$, otherwise $\rho' = \pi$.

Suppose $\rho' \neq \pi$. Let g be (a \mathbb{P} -name for) $f_i \cup \{(\text{FP}_{j-1}(f_i), \pi(j-1))\}$ and $q = q_\eta^i$. Then $q \geq q_{\eta \upharpoonright (i-1)}^{i-2}$, $q \Vdash f_{i-2} \leq_{i-2}^* g$, and

$$q \Vdash "g * (\rho' \upharpoonright (j-1)) = f_i * \rho' \text{ and } g * (\pi \upharpoonright (j-1)) = f_i * \pi".$$

Hence

$$(q, g * (\rho' \upharpoonright (j-1))) \Vdash " \tau \upharpoonright m_i = \sigma_{\rho', \eta}^i " \quad \text{and} \quad (q, g * (\pi \upharpoonright (j-1))) \Vdash " \tau \upharpoonright m_i = \sigma_{\pi, \eta}^i ".$$

Now we use our assumption that $\sigma_{\rho', \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \eta}^i \upharpoonright m_{i-2}$ (and (\boxtimes)₁₁) and (\boxtimes)₆ to conclude that $\sigma_{\rho', \eta}^i = \sigma_{\pi, \eta}^i$. Trivially the same conclusion holds if $\rho' = \pi$, so we have justified that

$$\left\{ \sigma_{\pi, \nu}^i : \nu \in \max(s_i) \ \& \ \pi \in {}^{[i/4]}n \ \& \ \sigma_{\rho, \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \nu}^i \upharpoonright m_{i-2} \right\} \subseteq \left\{ \sigma_{\pi, \eta}^i : \pi \in {}^j n \ \& \ \rho \upharpoonright (j-1) = \pi \upharpoonright (j-1) \right\}$$

and the latter set is of size at most n .

CASE B: $i = 4j + 3$ for some $j < \omega$.

Again, let us assume that $\eta, \nu \in \max(s_i)$, $\rho, \pi \in {}^{[i/4]}n$ are such that $\sigma_{\rho, \eta}^i \neq \sigma_{\pi, \nu}^i$ but $\sigma_{\rho, \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \nu}^i \upharpoonright m_{i-2}$. Then, like in the previous case, (\boxtimes)₇ implies $\eta \upharpoonright (i-1) = \nu \upharpoonright (i-1)$. Also $[i/4] = j = [(i-2)/4]$, so $\rho \upharpoonright [(i-2)/4] = \rho$, $\pi \upharpoonright [(i-2)/4] = \pi$. Now, if $\rho = \pi$, then trivially $\sigma_{\pi, \nu}^i = \sigma_{\rho, \nu}^i$. If $\rho \neq \pi$, then we use (\boxtimes)₆ (with $i-2, \rho, \pi, q_\eta^i, f_i$ here in place of i, ρ, ρ', q, g there, respectively) to argue that $\sigma_{\pi, \nu}^i = \sigma_{\rho, \nu}^i$. Consequently

$$\left\{ \sigma_{\pi, \nu}^i : \nu \in \max(s_i) \ \& \ \pi \in {}^{[i/4]}n \ \& \ \sigma_{\rho, \eta}^i \upharpoonright m_{i-2} = \sigma_{\pi, \nu}^i \upharpoonright m_{i-2} \right\} \subseteq \left\{ \sigma_{\rho, \nu}^i : \nu \in \max(s_i) \ \& \ \eta \upharpoonright (i-2) = \nu \upharpoonright (i-2) \right\}$$

and the latter set is of size at most n .

Now in both cases we easily get the assertion of (\boxtimes)₁₂, completing the proof of the theorem.

REFERENCES

- [1] Stefan Geschke. More on convexity numbers of closed sets in \mathbb{R}^n . *Proceedings of the American Mathematical Society*, 133:1307–1315, 2005.
- [2] Stefan Geschke and Menachem Kojman. Convexity numbers of closed sets in \mathbb{R}^n . *Proceedings of the American Mathematical Society*, 130:2871–2881, 2002.
- [3] Stefan Geschke, Menachem Kojman, Wiesław Kubiś, and Rene Schipperus. Convex decompositions in the plane and continuous pair colorings of the irrationals. *Israel Journal of Mathematics*, 131:285–317, 2002.
- [4] Thomas Jech. *Set theory*. Academic Press, New York, 1978.
- [5] Ludomir Newelski and Andrzej Roslanowski. The ideal determined by the unsymmetric game. *Proceedings of the American Mathematical Society*, 117:823–831, 1993.

- [6] Andrzej Rosłanowski. n -localization property. *Preprint*. math.LO/0507519.
- [7] Andrzej Rosłanowski. Mycielski ideals generated by uncountable systems. *Colloquium Mathematicum*, LXVI:187–200, 1994.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA AT OMAHA, OMAHA, NE 68182-0243, USA

E-mail address: `roslanow@member.ams.org`

URL: `http://www.unomaha.edu/logic`

DEPARTMENT OF MATHEMATICS, YORK UNIVERSITY, 4700 KEELE STREET, TORONTO, ONTARIO, CANADA M3J 1P3

E-mail address: `steprans@yorku.ca`

URL: `http://www.math.yorku.ca/~steprans`